

AUTHOR:

*Padeyev, L. D.*  
Padeyev, L. D.

20-5-8/54

TITLE:

On an Expression for the Trace of the Difference of Two Singular Differential Operators of the Sturm-Liouville's Type (O vyrazhenii dlya sleda raznosti dvukh singulyarnykh differentsial'nykh operatorov tipa Shturma-Liuvillya)

PERIODICAL: Doklady Akad.Nauk SSSR 1957, Vol.115, Nr.5, pp.878-881 (USSR)

ABSTRACT:

The formula for the trace of the difference of two regular Sturm-Liouville's operators with a discrete spectrum found by Gelfand and Levitan [Ref.1] is extended by the author in the simplest special case to operators with a continuous spectrum. The operators  $L_1 y = -y'' + q_1(x)y$ ,  $y(0) = 0$ ,  $i=1,2$ , are considered.

Theorem: If 1)  $\int_0^\infty x |q_1(x)| dx < \infty$ ,  $i=1,2$ , 2)  $g(x) = q_1(x) - q_2(x)$  is continuous in the neighborhood of  $x = 0$  and 3)  $\int_0^\infty g(x) dx = 0$ ,

then the trace of  $L_1 - L_2$  is finite and there holds the formula

Card 1/2

$$\text{Sp} (L_1 - L_2) = -\frac{1}{4} (q_1(0) - q_2(0)).$$

On an Expression for the Trace of the Difference of Two Singular  
Differential Operators of the Sturm-Liouville's Type (Cont.)

20-5-8/54

ASSOCIATION: Leningrad State University im. A. A. Zhdanov (Leningradskiy gosudar-  
stvennyy universitet im. A. A. Zhdanova)

PRESENTED: By V. I. Smirnov, Academician, March 1, 1957

SUBMITTED: February 20, 1957

AVAILABLE: Library of Congress

Card 2/2

LADYZHENSKAYA, O. A. and FADDEYEV, L. D.

"Perturbation Theory of a Continuous Spectrum."

paper submitted at International Congress Mathematicians, Edinburgh, 14 - 21 Aug  
1958.

24 (5)

AUTHOR: Faddeyev, L. D.

SOV/56-35-2-17/60

TITLE: On the Dispersion Relations in the Nonrelativistic Scattering Theory (O dispersionnykh sootnosheniyakh v nerelyativistskoy teorii rasseyaniya)

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1958, Vol 35, Nr 2, pp 433-439 (USSR)

ABSTRACT: The present paper is a continuation of several papers: (Wong, Khuri, Van Kampen, Jost, and Pais, references 1 - 4, - investigation of dispersion conditions with S-matrix at complex energies - with special causality principle - according to the theory developed by Fredholm (Fredgol'm)). In the present paper the author suggests a relatively simple derivation of dispersion conditions, in which the problem is reduced to an investigation of the properties of Green's (Grin) function of the total Hamiltonian. Derivation is of a general character and is suited for any amplitudes of scattering on a potential. The author demonstrates this method on the basis of the problem of scattering on a fixed three-dimensional center (nonrelativistic ansatz).

Card 1/2

On the Dispersion ~~Relations~~ in the Nonrelativistic  
Scattering Theory

SOV/56-35-2-17/60

In this way it is possible to verify the results obtained by Wong and Khuri. In the last chapter of the paper the connection between the dispersion relations and the problems relating to the complete characteristic of the S-matrix, (by means of the K-matrix  $S = (1 + iK)(1 - iK)^{-1}$ ) for  $0 \leq E < \infty$  is discussed for different scattering equations. There are 14 references, 7 of which are Soviet.

ASSOCIATION: Leningradskiy gosudarstvennyy universitet (Leningrad State University)

SUBMITTED: March 16, 1958

Card 2/2

AUTHOR: Ladyzhenskaya, O.A. and Faddeyev, L.D. SOV/20-120-6-5/59

TITLE: On the Perturbation Theory of the Continuous Spectrum (K teorii vozmushcheniy nepreryvnogo spektra)

PERIODICAL: Doklady Akademii nauk SSSR, Vol 120, Nr 6, pp 1187-1190 (USSR), 1958

ABSTRACT: Let  $K$  be an integral operator and let  $L_0$  denote the multiplication with the independent variable. The investigation of the spectrum of  $L = L_0 + \varepsilon K$  led Friedrichs [Ref 1,2] to the consideration of the integral equation

$$(1) \quad r(\lambda, \mu) = k(\lambda, \mu) + i\sqrt{\varepsilon} k(\lambda, \mu) r(\mu, \mu) + \varepsilon P \left( \frac{k(\lambda, \sigma) r(\sigma, \mu)}{\mu - \sigma} \right)$$

The solubility of (1) was proved by Friedrichs for small  $\varepsilon$  only. The authors prove that (1) is solvable for an arbitrary finite  $\varepsilon$ , and they present some properties of the spectrum of  $L$  without restriction to small  $\varepsilon$ . There are 2 non-Soviet references, 1 of which is German, and 1 American.

Card 1/2

On the Perturbation Theory of the Continuous Spectrum SOV/20-120-6-5/59

ASSOCIATION: Leningradskoye otdeleniye matematicheskogo instituta imeni  
V.A. Steklova (Leningrad Section of the Mathematical Institut  
imeni V.A. Steklov of the Academy of Sciences of the USSR)

PRESENTED: February 17, 1958, by V.I. Smirnov, Academician

SUBMITTED: February 10, 1958

1. Perturbation theory 2. Spectroscopy

Card 2/2

AUTHOR: Addeev, L. D. 87/20-121-1-10, '55

TITLE: On the Connection Between  $\delta$ -Matrix and Potential for a One-Dimensional Schrödinger Operator (O svyazi  $\delta$ -matritsy i potentsiala dlya odnomernogo operatora Shrelingera)

SYNOPSIS: Doklady Akademii nauk SSSR, 1958, Vol. 121, No. 1, pp. 63-66 (USSR)

ABSTRACT: The author investigates the one-dimensional Schrödinger equation 
$$Ly - d^2/dx^2 y + q(x)y = k^2 y$$
 on the whole axis  $-\infty < x < \infty$ , where the condition 
$$\int_{-\infty}^{\infty} (1 + |x|) |q(x)| dx < \infty$$
 is assumed to be satisfied.

If this condition is satisfied the solution  $f_1(x, k)$  (for which  $\lim_{x \rightarrow \infty} e^{-ikx} f_1(x, k) = 1$

Card 1/4 holds) and  $f_2(x, k)$  (for which  $\lim_{x \rightarrow -\infty} e^{ikx} f_2(x, k) = 1$  holds)



SOV/20-121-1-16/55

On the Connection Between S-Matrix and Potential for a One-Dimensional Schrödinger Operator

exist for every  $k$  of the upper semiplane. In  $k > 0$ . According to B. Ya. Levin a representation

$$f_1(x, k) = e^{ikx} + \int_x^\infty A_1(x, y) e^{iky} dy \text{ exists, whereby}$$

$$\int_a^\infty dx \int_x^\infty dy |A_1(x, y)|^2 \leq C_a, \quad a > -\infty. \text{ Furthermore is valid}$$

$$f_2(x, k) = e^{-ikx} + \int_{-\infty}^x A_2(x, y) e^{-iky} dy \quad \int_{-\infty}^b dx \int_{-\infty}^x dy |A_2(x, y)|^2 \leq C_b,$$

$b < \infty$ . In the case of real  $k \neq 0$  the solutions  $f_1(x, k)$  and  $f_1(x, -k) = f_1(x, k)$  or  $f_2(x, k)$  and  $f_2(x, -k) = f_2(x, k)$  form a complete system thus that every solution can be represented

Card 2/4

On the Connection Between S-Matrix and Potential for a One-Dimensional Schrödinger Operator

SOV/20-121-1-16/55

as their linear combination.

$$f_1(x, k) = c_{11}(k) f_1(x, k) + c_{12}(k) f_1(x, -k)$$

$$f_1(x, k) = c_{22}(k) f_2(x, k) + c_{12}(k) f_2(x, -k) \text{ holds in particular.}$$

A lemma on the determination of these coefficients and other lemma are given. The equations derived in this paper allow to ascertain the potential by means of the S-matrix. The author wants to find out which properties the elements of the S-matrix must exhibit that the potential satisfies the condition

$$\int_{-\infty}^{\infty} (1 + |x|) q(x) dx < \infty. \text{ The inverse problem}$$

-∞

can be solved by means of a lemma given in this paper. The result is formulated as a theorem. There are 6 references, 5 of which are Soviet.

Card 3/4

007/20-121-1-16/55

On the Connection Between S-Matrix and Potential for a One-Dimensional  
Schrödinger Operator

ASSOCIATION: Leningradskiy gosudarstvennyy universitet im. A. A. Zhdanova  
(Leningrad State University imeni A. A. Zhdanov)

PRESENTED: March 31, 1958, by V. I. Smirnov, Member, Academy of Sciences.  
NOTE (this date is obviously a printing error in the Soviet  
paper)

SUBMITTED: March 12, 1958

1. Operators (Mathematics) 2. Mathematics

FADDEYEV, L. D. Cand Phys-Math Sci -- (diss) "Properties of <sup>an matrix</sup> ~~the~~ S-shaped  
stamper for ~~discussion~~ on a local potential." Len, 1959. 8 pp (Len Order of  
Lenin State Univ im A. A. Zhdanov), 150 copies (KL, 45-59, 143)

16(1),16(2),24(5)

AUTHOR: Faddeyev, L.D.

SOV/42-14-4-3/27

TITLE: The Reversion Problem of the Quantum Theory of Scattering

PERIODICAL: Uspekhi matematicheskikh nauk, 1959, Vol 14,Nr 4,pp 57-120 (USSR)

ABSTRACT: This is a connected detailed representation of the results obtained during the last 12 years by the investigation of the reversion problem of the quantum theory which is formulated by the author as follows: Under the assumption that the potential  $q(x)$  for  $x \rightarrow \infty$  decreases sufficiently fast. let the solution of the equation

$$(0.1) \quad L\psi = -\frac{d^2}{dx^2} \psi(x,k) + q(x) \psi(x,k) = k^2 \psi(x,k)$$

$$(0.2) \quad \psi(0,k) = 0$$

have the asymptotic

$$(0.3) \quad \psi(x,k) \approx \mathcal{C}(k) \sin(kx - \eta(k)).$$

How far is  $q(x)$  determined by  $\eta(k)$ ? How are the properties of  $q(x)$  and  $\eta(k)$  combined with each other?

By a skilful combination of the methods of I.M.Gel'fand and B.M.Levitan on the one hand, and by V.A.Marchenko and M.G.Kreyn

Card 1/2

The Reversion Problem of the Quantum Theory of  
Scattering

SOV/42-14-4-3/27

at the other hand, the author obtains a clear theory. The most difficult mathematical proofs are replaced by physically obvious considerations. The paper contains 15 paragraphs, the first 13 of which are devoted to the case (0.1)-(0.3), while the last two paragraphs treat the case

$$-\frac{d^2}{dx^2} \psi(x,k) = \left( \frac{l(l+1)}{x^2} + q(x) \right) \psi(x,k) = k^2 \psi(x,k)$$

with  $l > 0$ .

Beside of the above Soviet scientists the author mentions: O.A.Ladyzhenskaya, Z.S.Agranovich, A.Ya.Povzner, B.Ya. Levin, L.A.Chudov, M.G.Neygauz V.V.Stashevskaya, V.Ya. Volk, and A.Sh. Blokh.

There are 66 references, 25 of which are Soviet, 21 American, 5 German, 3 Danish, 9 Italian, 2 Swiss, and 1 Norwegian.

SUBMITTED: February 12, 1959

Card 2/2

00924

S/056/60/039/005/041/051  
B006/B077

24.450

AUTHOR:

Faddeyev, L. D.

TITLE:

Scattering Theory for a Three-particle System

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1960,  
Vol. 39, No. 5(11), pp. 1459 - 1467

TEXT: The equations presently obtained in the scattering theory for the Hamiltonian eigenfunctions of the system in the configuration space show a number of shortcomings such as the Lippman-Schwinger integral equations. The author proposes equations for the eigenfunctions which do not have these shortcomings. It can be shown that the eigenfunctions of the Hamiltonian for a three-particle system with pair interaction can be easily represented as the sum of three terms. There is a linked set of equations for each of these terms. All equations are inhomogeneous; only for energies corresponding to a bound state of the system exists a solution in form of a homogeneous equation. In order to determine the kernels of the integral equation only the pair problems are to be solved. These kernels are found to be generalizations of the so-called

Card 1/3

86924

Scattering Theory for a Three-particle System S/056/60/039/005/041/051  
B006/B077

T-matrix and are easily to be determined for various limiting cases, if no potential exists. The separation of the wave function into three parts has been done previously (in three-body problems) very successfully (Refs. 7,8). In the first part of this study the equations are derived formally for a system consisting of three nonrelativistic spin-zero particles with different mass. Two of these particles are assumed to be in a bound state. The second part of this work deals with the transformation of the obtained equations into momentum representation and the meaning of the model of paired particles (nuclei) is investigated. The last part gives the discussions of the results. It is shown that in opposition to the Lippman-Schwinger type equations these equations have a unique solution. If the limit is crossed towards vanishing interaction range, the well known equations by G. V. Skornyakov and K. A. Ter-Martirosyan are obtained. The author thanks F. A. Berezin, V. N. Gribov, S. V. Maleyev, and L. V. Prokhorov for discussions. There are 10 references: 2 Soviet and 8 US.

Card 2/3



86924

Scattering Theory for a Three-particle System S/056/60/039/005/041/051  
B006/B077

ASSOCIATION: Matematicheskii institut Akademii nauk SSSR (Mathematics  
Institute of the Academy of Sciences USSR)

SUBMITTED: July 30, 1960

Card 3/3

80035

S/020/60/132/01/02/064

16.3400 16.4600

AUTHORS: Buslayev, V.S., and Faddeyev, L.D.

TITLE: Formulas for Traces in the Case of Sturm - Liouville's Differential Singular Operator

PERIODICAL: Doklady Akademii nauk SSSR, 1960, Vol. 132, No. 1, pp. 13-16

TEXT: The authors consider the operator  $Ly = -y'' + q(x)y$ ,  $y(0) = 0$ , where

it is assumed that  $\int_0^{\infty} x|q(x)| dx < \infty$ . The spectrum of  $L$  consists of the

continuable part  $[0, \infty]$  and finitely many negative values  $\lambda_1 = -\alpha_1^2$  ( $\alpha_1 > 0$ ;  $1 = 1, 2, \dots, m$ ). Let

$$M(s) = 1 + \int_0^{\infty} e^{isx} q(x) \varphi(x, s) dx = \Lambda(s) e^{i\eta(s)}$$

$$(s = \sigma + i\tau, 0 \leq \tau < \infty, -\infty < \sigma < \infty)$$

(for the notations see (Ref. 2)). Let  $R_{\lambda}$  be the resolvent of  $L$ ; the upper index 0 relates to the case  $q(x) = 0$ .  $\lambda_1$  ( $1 = 1, 2, \dots, m$ ) it holds

Card 1/4

Formulas for Traces in the Case of Sturm -  
Liouville's Differential Singular Operator

80035  
S/020/60/132/01/02/064

$$\text{Sp}(R_\lambda - R_\lambda^0) = - \frac{d}{d\lambda} \ln M(\sqrt{\lambda}) ; 0 \leq \arg \sqrt{\lambda} \leq \pi .$$

Conclusion :  $M(\sqrt{\lambda}) = \det (E + q R_\lambda^0) .$

For  $q(x) \in L[0, \infty]$  it holds

$$(\alpha) \quad \text{Sp}(R_\lambda - R_\lambda^0) = - \int_{-\infty}^{\infty} \xi(t) d \frac{1}{t - \lambda} ,$$

where

$$\xi(t) = \begin{cases} \frac{1}{\pi} \tau(\sqrt{t}) & t > 0 \\ - \int_{-\infty}^t \sum_1 \delta(z - \lambda_1) dz , & t < 0 . \end{cases}$$

Let for  $x \geq 0$  exist the continuous  $q^{(n)}(x)$  ( $n \geq 1$ ) ; where  $\lim_{x \rightarrow \infty} q^{(1)}(x) = 0$

for  $l = 0, \dots, n$  . Let further

Card 2/ 4

4

80035

Formulas for Traces in the Case of Sturm -  
Liouville's Differential Singular Operator

S/020/60/132/01/02/064

$$V_1 = \lim_{\alpha \rightarrow \infty} V_1(\alpha) ; V_0(\alpha) = - \int_0^\alpha q(z) dz ,$$

$$V_1(\alpha) = q^{(1-1)}(0) + \sum_{m=0}^{l-1} C_{1-1}^m \int_0^\alpha dz V_m(z) q^{(1-m-1)}(z)$$

$$Q_p = V_{p-1} + \sum_{j=1}^{p-1} \frac{1}{p} V_{p-j-1} Q_j$$

Theorem 2 : Under the given assumptions it holds

$$\begin{aligned} (-1) \sum_{l=1}^m x_l^{2\mu} + \frac{2\mu}{\pi} \int_0^\infty k^{2\mu-1} \left[ \eta(k) - \sum_{l=0}^{\mu-1} \frac{(-1)^{l+1}}{(2k)^{2l+1}} Q_{2l+1} \right] dk = \\ = (-1)^\mu \frac{\mu}{2^{2\mu}} Q_{2\mu} \quad (\mu = 1, 2, \dots \leq \frac{n}{2}) ; \end{aligned}$$

Card 3/4

Formulas for Traces in the Case of Sturm -  
Liouville's Differential Singular Operator

80035  
S/020/60/132/01/02/064

$$(-1)^\mu \sum_{l=1}^m a_1^{2\mu+1} - \frac{2\mu+1}{\pi} \int_0^\infty k^{2\mu} \left[ \ln A(k) - \sum_{l=1}^{\mu} \frac{(-1)^{l+1}}{(2k)^{2l}} Q_{2l} \right] dk =$$

$$= (-1)^\mu \frac{2\mu+1}{2^{2\mu+2}} Q_{2\mu+1} \quad (\mu = 0, \dots, \leq \frac{n-1}{2}) .$$

The authors mention I.M. Gel'fand, B.M. Levitan, L.A. Dikiy and I.M. Lifshits.  
The authors thank M.G. Kreyn and M.Sh. Birman for discussions.  
There are 7 references : 6 Soviet and 1 American.

ASSOCIATION: Leningradskiy gosudarstvennyy universitet imeni A.A. Zhdanova  
(Leningrad State University imeni A.A. Zhdanov)

PRESENTED: January 3, 1960, by V.I. Smirnov, Academician

SUBMITTED: December 17, 1959

Card 4/4

4

S/056/61/041/006/029/054  
B146/B102

AUTHORS: Minlos, R. A., Faddeyev, L. D.

TITLE: The three-particle problem with point interaction

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 41,  
no. 6, 1961, 1850-1851

TEXT: An integral equation derived by K. A. Ter-Martirosyan and G. V. Skornyyakov (Ref. 1: ZhETF, 31, 775, 1956) for the wave function of a three-particle system with point interaction is considered on the basis of scalar and homogeneous particles. Besides a relation following from the asymptotic behavior and from the orthogonality of the solutions, this equation has solutions corresponding to an infinite set of bound states. The proof furnished for this is only applicable to spherically symmetric solutions. The work is based on the Ter-Martirosyan - Skornyyakov model improved by G. S. Danilov (Ref. 2: ZhETF, 40, 498, 1961); a more general treatment in a mathematical paper by the authors (Ref. 4: R. A. Minlos, L. D. Faddeyev, DAN SSSR, 141, 6, 1961) is referred to. There are 4 Soviet references.

Card 1/2

The three-particle problem with ...

S/056/61/041/006/029/054  
B146/B102

ASSOCIATION: Moskovskiy gosudarstvennyy universitet (Moscow State  
University)

SUBMITTED: June 16, 1961



Card 2/2

21959

S/020/61/137/005/002/026  
C111/C222

21.4500

AUTHORS: Berezin, F.A., and Faddeyev, L.D.

TITLE: A remark on the Schrödinger equation with a singular potential

PERIODICAL: Akademiya nauk SSSR. Doklady, vol.137,no.5,1961, 1011-1014

TEXT: The solution of the equation

$$-\Delta \psi + \varepsilon \delta(x) \psi = E \psi, \quad (1)$$

where  $\delta(x)$  is the Dirac  $\delta$ -function, contains certain difficulties since

$$H = -\Delta + \varepsilon \delta(x) \quad (2)$$

is no operator in the Hilbert space.

Instead of (1) the authors consider

$$-\Delta \psi + \varepsilon(N) \int u_N(x,y) \psi(y) d^3y = E \psi, \quad (N)$$

where  $u_N$  has the property

$$\lim_{N \rightarrow \infty} u_N(x,y) = \delta(x) \delta(y). \quad (3)$$

For the solution of (N) the authors use the Fourier transformation and obtain

Card 1/5



21959

A remark on the Schrödinger equation...

3/020/61/137/005/002/026  
C111/C222

$$p^2 \tilde{\psi} + \frac{(N)}{8\pi^3} \int \tilde{u}_N(p,q) \tilde{\psi}(q) d^3q = E \tilde{\psi},$$

$$\tilde{u}_N(p,q) = \int e^{i(qy-px)} u_N(x,y) d^3x d^3y, \quad (\tilde{N})$$

where

$$\lim_{N \rightarrow \infty} \tilde{u}_N(p,q) = 1. \quad (3')$$

Now  $u_N$  is chosen so that

$$\tilde{u}_N(p,q) = \chi_N(p) \chi_N(q), \quad \chi_N(p) = \begin{cases} 1 & \text{for } p^2 < N^2, \\ 0 & \text{for } p^2 > N^2. \end{cases} \quad (4)$$

Then the eigenfunctions belonging to the continuous spectrum read

Card 2/5

A remark on the Schrödinger equation...

S/020/61/137/005/002/026  
C111/C222

$$\tilde{\psi}_N^+(p, s) = \delta(p-s) - \frac{\varepsilon'(N)}{1 + \varepsilon'(N) \int \frac{\chi_N^2(p) d^2p}{p^2 - s^2 - i0}} \frac{\chi_N(p) \chi_N(s)}{p^2 - s^2 - i0}. \quad (5)$$

$$s^2 = E, \quad \varepsilon' = \frac{\varepsilon(N)}{8\pi^2}.$$

Furthermore:

$$\int \frac{\chi_N^2(p) d^2p}{p^2 - s^2 - i0} = 4\pi \int_0^N \frac{p^2 dp}{p^2 - s^2 - i0} = 4\pi \left( N + \frac{|s|}{2} (-\pi i + \ln \frac{N - |s|}{N + |s|}) \right). \quad (6)$$

Choosing  $\varepsilon'(N) = \frac{\alpha}{1 - 2\pi\alpha N}$ ,  $\alpha = \text{const}$ , then the limit value of  $\tilde{\psi}_N^+$  for  $N \rightarrow \infty$  equals

$$\tilde{\psi}^+ = \delta(p-s) - \frac{\alpha}{1 - 2\pi^2 i \alpha |s|} \frac{1}{p^2 - s^2 - i0}. \quad (7)$$

At the other hand, the authors consider the Fourier transform of (2)

$$\tilde{H}\psi = p^2\psi + \varepsilon' \int \psi d^3p. \quad (8)$$

Card 3/5

219yy

S/020/61/137/005/002/026

C111/C222

A remark on the Schrödinger equation...

Let  $D_L$  be the set of functions for which

$$\int p^4 |\psi|^2 d^3p < \infty, \quad \int \psi d^3p = 0.$$

Let  $L$  be the operator of the multiplication with  $p^2$  defined in  $D_L$ .  $L$  is a closed symmetrical operator with the defect index  $(1,1)$ . All extensions of  $L$  are given by

$$H_\alpha \psi = p^2 \psi + \lim_{N \rightarrow \infty} \frac{\alpha}{1-4\pi\alpha N} \int \chi_N(p) \psi(p) d^3p, \quad (9)$$

where  $\psi(p)$  has the properties

$$\int \chi_N(p) \psi(p) d^3p = c(1-4\pi\alpha N) + o(1), \quad \int |H_\alpha \psi|^2 d^3p < \infty. \quad (9')$$

It is stated that the eigenfunctions of the continuous spectrum  $H_\alpha$  are given by (7). Using these results then the scattering operator and the results given in (Ref.1: Ya.B.Zel'dovich, Zh E T F 38, no.3, 819(1960)) can be obtained.

In the  $x$ -representation it holds

$$H_\alpha f = -\Delta f + \alpha \lim_{N \rightarrow \infty} \frac{1}{1-4\pi\alpha N} \frac{\sin N|x|}{|x|} \int \frac{\sin N|y|}{|y|} f(y) d^3y. \quad (12)$$

Card 4/5

21959

S/020/61/137/005/002/026  
C111/C222

A remark on the Schrödinger equation...

The region of definition of  $H_\alpha$  consists of functions satisfying the condition

$$\int \frac{\sin N|x|}{|x|} f(x) d^3x = o(1-4\pi\epsilon N) + o(1), \quad \int |H_\alpha f|^2 d^3x < \infty. \quad (12')$$

It is stated that the mathematical content of the investigation of the equation (1) by physicists (e.g. Ref.1) consists in replacing (2) by the operator (12'), (12') being an extension of the operator  $-\Delta$  the region of definition of which consists of those  $f(x)$  for which  $f(0)=0$ .

There are 2 Soviet-bloc references.

PRESENTED: November 25, 1960, by I.G.Petrovskiy, Academician

SUBMITTED: November 24, 1960

Card 5/5

24037

S/020/61/138/003/009/017  
B104/B205

24.4400

AUTHOR: Faddeyev, L. D.

TITLE: Construction of the resolvent of the Schroedinger operator of a system of three particles with pair interaction

PERIODICAL: Doklady Akademii nauk SSSR, v. 138, no. 3, 1961, 565 - 567

TEXT: The energy operator for a system of N particles having the masses  $m_1, \dots, m_N$ , which undergo pair interaction, has the form

$$H_N = - \sum_{i=1}^N \frac{1}{2m_i} \nabla_i^2 + \sum_{i<j}^N v_{ij}(r_i - r_j). \quad (1)$$

Only the operator  $H_2$  has been studied so far. In this connection, the author refers to papers by A. Ya. Povzner (Matem. sborn., 32, 1, (1953); DAN, 104, no. 3, 360 (1955)), T. Kato (Comm. Pure and Appl. Math., 12, 403 (1959)), and T. Ikebe (Arch. Rat. Mech. Anal., 5, no. 1, 1(1960)). In a previous paper (ZhETF, 39, no. 11, 1569 (1960)), the author suggested a new integral equation for studying a system of three particles. In the present paper, he reports the results of a study on the behavior of the

Card 1/6

24037

S/020/61/138/003/009/017

B104/B205

Construction of the resolvent...

resolvent of the operator  $H_3$ , which he obtained from this equation. Instead of  $H_3$ , the operator  $H$  is studied, which was obtained from  $H_3$  by proceeding to pulse representation, i. e., after a Fourier transformation, and by separating the operator from the kinetic energy of the center of mass. Here, it is assumed that  $m_1 = m_2 = m_3$ . The three vectors assumed,  $p_1$ ,  $p_2$ , and  $p_3$ , are given by

$$p_1 + p_2 + p_3 = 0 \quad (2).$$

Each pair of these vectors traverses the six-dimensional space  $E_6$  independently. The operator  $H$  is given in the  $\mathcal{L}_2(E_6)$  space, and has the form

$$H = H_0 + V = H_0 + V_{23} + V_{31} + V_{12} \quad (3),$$

where  $H_0$  is an operator acting on the function

$$p_1^2 + (p_1 p_2) + p_2^2 = p_2^2 + (p_2 p_3) + p_3^2 = p_3^2 + (p_3 p_1) + p_1^2 \quad (4).$$

The operator  $V_{23}$  has the kernel

$$V_{23}(p, p') = v_{23}(p_2 - p_2') \delta(p_1 - p_1') \quad (5).$$

Card 2/6

24037

S/020/61/138/003/009/017

B104/B205

Construction of the resolvent...

The operators  $V_{31}$  and  $V_{12}$  are defined analogously. If  $R(z) = (H - zI)^{-1}$  and  $R_0(z) = (H_0 - zI)^{-1}$  are the resolvents of the operators  $H$  and  $H_0$ , one has

$$R(z) = R_0(z) - R_0(z)VR(z). \quad (7)$$

If  $R(z)$  is sought in the form

$$R(z) = R_0(z) - R_0(z)T(z)R_0(z). \quad (8)$$

then  $T(z)$  is given by

$$T(z) = V - VR_0(z)T(z) \quad (9).$$

Eqs. (7) and (9) are particularly valuable for studying the operator  $H_2$ . The operator equation

$$X(z) = \begin{pmatrix} V_{33} & 0 & 0 \\ 0 & V_{31} & 0 \\ 0 & 0 & V_{11} \end{pmatrix} - \begin{pmatrix} V_{33} & V_{32} & V_{30} \\ V_{31} & V_{31} & V_{31} \\ V_{11} & V_{11} & V_{11} \end{pmatrix} R_0(z)X(z). \quad (10)$$

is formulated in accordance with (9).  $T_{23}(z)$  is supposed to be a solution to the equation

$$T_{23}(z) = V_{23} - V_{23}R_0(z)T_{23}(z) \quad (11),$$

Card 3/6

24037

S/020/61/138/003/009/017  
B104/B205

Construction of the resolvent...

which also holds for  $T_{31}(z)$  and  $T_{12}(z)$ . Thus, the following relation is obtained from Eq. (10):

$$\begin{aligned} \mathcal{X}(z) = & \begin{pmatrix} T_{33}(z) & 0 & 0 \\ 0 & T_{11}(z) & 0 \\ 0 & 0 & T_{11}(z) \end{pmatrix} - \begin{pmatrix} 0 & T_{33}(z) & T_{33}(z) \\ T_{11}(z) & 0 & T_{11}(z) \\ T_{11}(z) & T_{11}(z) & 0 \end{pmatrix} R_0(z) \mathcal{X}(z) = \\ & = \mathcal{X}_0(z) - \mathcal{M}(z) \mathcal{X}(z). \end{aligned} \quad (12)$$

It is easily seen that  $T_{23}(p, p') = t_{23}(-p_2 - \frac{1}{2}p, -p_2' - \frac{1}{2}p_1,$

$$z - \frac{3}{4}p_1^2) \delta(p_1 - p_1') \quad (13),$$

where  $t_{23}$  is the solution to the integral equation

$$t(k, k', z) = v(k - k') - \int v(k - k'') (k''^2 - z)^{-1} t(k'', k', z) dk'' \quad (14).$$

In order to find out whether there are no  $\delta$ -singularities in the free term, the matrix  $\mathcal{M}(z) = \mathcal{X}(z) - \mathcal{X}_0(z)$  is studied, for which the following relations are valid:

Card 4/6



24037

S/020/61/138/003/009/017

B104/B205

Construction of the resolvent...

where  $\mathbb{M}(z) = \mathbb{M}_0(z) - \mathbb{M}(z) \mathbb{M}(z), \quad (15)$

$$\mathbb{M}_0(z) = \begin{pmatrix} 0 & T_{12}(z) R_0(z) T_{11}(z) & T_{12}(z) R_0(z) T_{11}(z) \\ T_{11}(z) R_0(z) T_{12}(z) & 0 & T_{11}(z) R_0(z) T_{11}(z) \\ T_{11}(z) R_0(z) T_{12}(z) & T_{11}(z) R_0(z) T_{11}(z) & 0 \end{pmatrix}. \quad (16)$$

Using Eq. (15) it is possible to investigate the behavior of  $R(z)$  for  $\text{Im } z \neq 0$ . The following theorem has been established: If the potential  $V_{ij}$  satisfies the condition

$$|v_{ij}(q)| \leq C(1+|q|)^{-1-\epsilon}, \quad \epsilon_0 > 0, \quad i, j = 1, 2, 3. \quad (6)$$

the resolvent of the operator  $H$  will acquire the form

$$R(z) = R_0(z) + \sum_{j=1}^3 (R_{1j}(z) - R_0(z)) + R_0(z) W(z) R_0(z) \quad (22).$$

Here,  $R_0(z)$  and  $R_{1j}(z)$  are the resolvents of the operators  $H_0, \dots, H_{1j}$ , where  $H_{1j} = H_0 + V_{1j}$  ( $i, j = 1, 2, 3$ );  $W(z)$  is an integral operator,

Card 5/6

24037

S/020/61/138/003/009/017  
B104/B205

Construction of the resolvent...

the kernel of which is given by

$$|W(p, p')| \leq C \sum_{j=1}^2 M_{1j}(p, p', \varepsilon) \quad (23),$$

where  $\varepsilon < \varepsilon_0$ . S. M. Nikol'skiy is mentioned. There are 6 references:  
4 Soviet-bloc and 2 non-Soviet-bloc.

ASSOCIATION: Leningradskoye otdeleniye Matematicheskogo instituta im.  
V. A. Steklova Akademii nauk SSSR (Leningrad Department of  
the Institute of Mathematics imeni V. A. Steklov of the  
Academy of Sciences USSR)

PRESENTED: January 30, 1961, by V. I. Smirnov, Academician

SUBMITTED: January 19, 1961

Card 6/6

MINIOS, R.A.; FADDEYEV, L.D.

Point interaction for a system of three particles in quantum mechanics.  
Dokl. AN SSSR 141 no.6:1335-1338 D '61. (MIRA 14:12)

1. Moskvoskiy gosudarstvennyy universitet im. M.V.Lomonosova.  
Predstavleno akademikom I.G.Petrovskim.  
(Operators (Mathematics)) (Quantum theory)

FADDEYEV, L. D.

"Construction of the resolvent of the energy operator for a three  
particle system and the scattering problem"

report submitted at the Intl Conf of Mathematics, Stockholm, Sweden,  
15-22 Aug 62

FADDEYEV, L.D.

Structure of the resolvent of Schrödinger's operator for a system of three particles, and the scattering problem. Dokl. AN SSSR 145 no.2:301-304 J1 '62. (MIRA 15:7)

1. Leningradskoye otdeleniye Matematicheskogo instituta imeni V.A. Steklova AN SSSR. Predstavleno akademikom V.I. Smirnovym. (Problem of three bodies) (Operators (Mathematics))

FADDEYEV, L.D.; PETROVSKIY, I.G., akademik, otv. red.; NIKOL'SKIY,  
S.M., prof., zam. otv. red.; TRAVIN, N.V., red. izd-va;  
SMIRNOVA, A.V., tekhn. red.

[Mathematical problems in the quantum theory of scattering  
for a system of three particles]. Matematicheskiy voprosy  
kvantovoi teorii rasseianiia dlia sistemy trekh chastits.  
Moskva, Izd-vo Akad. nauk SSSR, 1963. 119 p. (Akademiia nauk  
SSSR, Matematicheskii institut. Trudy, no.69).

(MIRA 16:4)

(Quantum theory) (Scattering(Physics))

FADDEYEV, L.D.

Separation of self-action and scattering effects under the  
perturbation theory. Dokl. AN SSSR 152 no.3:573-576 S '63.  
(MIRA 10:12)

1. Leningradskoye otdeleniye Matematicheskogo instituta im. V.A.  
Steklova AN SSSR. Predstavleno akademikom V.I.Smirnovym.

L 45909-65 EWT(d)/T IJP(c)

ACCESSION NR AM4043734

BOOK EXPLOITATION

S/ 30  
B/H

Vilenkin, N. YA.; Gorin, YE. A.; Kostyuchenko, A. G.; Kraonozel'skiy, M. A.;  
Kreyn, S. G.; Maslov, V. P.; Mityagin, B. S.; Potapin, IR. I.; Rutitskiy,  
YA. D.; Sobolev, V. I.; Stetsenko, V. YA.; Faddeyev, L. D.; Tsitlandze, E. S.

Functional analysis (Funktional'nyy analiz), Moscow, Izd-vo "Nauka", 1964,  
424 p. biblio., index. Errata slip inserted. 17,500 copies printed. Series  
note: Spravochnaya matematicheskaya biblioteka.

TOPIC TAGS: functional analysis, mathematics, operator equation, quantum  
mechanics, Hilbert space, Banach space, linear differential equation

PURPOSE AND COVERAGE: This issue in a series of Handbooks of the Mathematical  
Library contains much material grouped basically around the theory of  
operators and operator equations. It presents the basic concepts and methods  
of functional analysis, theory of operators in Hilbert space and in conical  
space, the theory of nonlinear operator equations, the theory of standard rings  
applied to equations in partial derivatives, to integral equations. A  
separate chapter is devoted to the basic operator of quantum mechanics. Citing  
of the theory of generalized functions takes up a large part of the book. The  
book explains mathematical facts; theorems and formulas, as a rule, are given

Card 1/2



L 45807-65

ACCESSION NR AM4043734

without proofs. Main attention is given to concepts without excessive detail. The book is intended for mathematicians, mechanical engineers, and physicists. It contains much of value for students and graduate students.

TABLE OF CONTENTS [abridged]:

Foreword	— 13
Ch. I. Basic concepts of functional analysis	— 17
Ch. II. Linear operators in Hilbert space	— 79
Ch. III. Linear differential equations in Banach space	— 146
Ch. IV. Nonlinear operator equations	— 187
Ch. V. Operators in space with a cone	— 229
Ch. VI. Commutative standard rings	— 256
Ch. VII. Quantum mechanics operators	— 279
Ch. VIII. Generalized functions	— 323
Bibliography	— 414
Subject Index	— 418

SUBMITTED: 06Feb64

SUB CODE: MA

NO REF SOV: 038

OTHER: 012

Card 2/2 01/

L 63360-65 EWT(d) IJP(c)

ACCESSION NR: AT5018144

UR/2517/64/073/000/0292/0313

AUTHOR: Faddeyev, L. D.

TITLE: On Friedrichs' model in the theory of perturbations of the continuous spectrum

SOURCE: AN SSSR. Matematicheskii institut. Trudy, v. 73, 1954. Krayevyye zadachi matematicheskoy fiziki (Boundary value problems in mathematical physics); sbornik rabot, no. 2, 292-313

TOPIC TAGS: continuous spectrum, quantum theory, Schroedinger equation, perturbation theory, integral operator, integral equation

ABSTRACT: Complete proofs are given for prior formulations made by the author and O. A. Ladyzhenskaya, who extended the work of Friedrichs and Povzner on perturbations of a continuous spectrum by showing that one may remove the limitations on the smallness of perturbations in Friedrichs' theory by assuming that the kernel of the operator is a completely continuous operator and that the Hölder index of the kernel is greater than  $\frac{1}{2}$ . The method used, however, differs somewhat from that used previously and was developed in work on the scattering problem for a system

Card 1/2

L 63360-65

ACCESSION NR: AT5018144

of three particles. The major analytic part of the work is devoted to the study of the integral equation:

$$t(x, y, \lambda) = v(x, y) - \int v(x, z) t(z, y, \lambda) (z - \lambda)^{-1} dz$$

for a  $t$ -matrix-kernel. Validity theorems are proved for convergence estimates for the  $T(\lambda)$  operator. A theorem for expansion with the operator is stated and proved.

Orig. art. has: 175 formulas

ASSOCIATION: none

SUBMITTED: 00

ENCL: 00

SUB CODE: GP, MA

NO REF SOV: 005

OTHER: 006

Card 2/2

FADDEYEV, L.D.

Properties of the S-matrix of a one-dimensional Schrodinger equation. Trudy Mat. Inst. 73:314-336 '64.

Friedrichs's model in the theory of perturbations of a continuous spectrum. Ibid.:292-313 (MIRA 18:3)

L 13486-65 ENT(1) LJP(c)/SSD/AS(mp)-2/AFWL/ESD(gs)/ESD(e)  
ACCESSION NR: AP4047899 S/0056/64/047/004/1315/1321

AUTHORS: Popov, V. N.; Faddeyev, L. D.

TITLE: Concerning one approach in the Bose gas theory at low temperatures

SOURCE: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 47, no. 4, 1964, 1315-1321

TOPIC TAGS: Bose Einstein gas, low temperature research, perturbation theory, annihilation, Green function, phonon

ABSTRACT: An approach is suggested for the theory of the Bose gas, believed to be more rigorous and simpler conceptually than the earlier treatments. The method is based on the premise that the existence of a condensate at low temperatures precludes the application of ordinary perturbation theory in which the unperturbed Hamiltonian is that of noninteracting particles, since in the ordinary theory

Card 1/3

L 13486-65

ACCESSION NR: AP4047899

the Green's function acquires nonphysical singularities when the temperature is reduced at fixed density. Consequently the authors start with a different unperturbed Hamiltonian which is made quadratic in the annihilation and creation operators by means of a specific canonical transformation. This perturbation theory is developed in diagram form, and the consequences of the resultant formalism is then discussed. In particular, it is shown how the phase transition associated with the appearance of the condensate shows up as the point where the canonical transformation degenerates into the identity transformation and the specially developed perturbation theory goes over into the usual one. The one-particle excitation spectrum is shown to have a phonon character below the transition temperature. Orig. art. has: 3 figures and 5 formulas.

ASSOCIATION: Leningradskoye otdeleniye Matematicheskogo instituta  
im. V. A. Steklova (Leningrad Division, Mathematics Institute,  
Academy of Sciences, USSR)

Card 2/3

L 13486-65

ACCESSION NR: AP4047899

SUBMITTED: 21Jan64

ENCL: 00

SUB CODE: GP, MA

NR REF SOV: 004

OTHER: 002

Card 3/3

FADDEYEV, L.D.

Growing solutions to the Schrödinger equation. Dokl. AN SSSR  
165 no.3:514-517 N '65. (MIRA 18:11)

1. Leningradskoye otdeleniye Matematicheskogo instituta im.  
V.A. Staklova AN SSSR. Submitted April 5, 1965.



FADDEYEV, M., inzh.

Modeling regulating structures and cuts on aerodynamic  
pressure models. Rech. transp. 20 no.12:44-45 D '61.  
(MIRA 14:12)

(Rivers—Regulation)  
(Aerodynamic models)

VERKHUNOV, P.M., kand.sel'skokhoz.nauk; FADDEYEV, M.G., mladshiy nauchnyy  
sotrudnik

Estimating the remnants of wood during the inspection of cutting  
places in heavily wooded areas of Siberia. Trudy VSNIPILesdrev  
no.5:11-14 '62. (MIRA 16:5)

1. Laboratoriya lesosyr'yevykh resursov Vostochno-Sibirskogo  
nauchno-issledovatel'skogo i proyektnogo instituta lesnoy i  
derevoobrabatyvayushchey promyshlennosti (for Verkhunov).  
(Siberia--Forest management)

FADDEYEV, M.P.

Problems of similitude related to aerodynamic stream bed models.  
Izv.Sib.otd.AN SSSR no.9:44-51 '60. (MIRA 13:11)

1. Novosibirskiy institut inzhenerov vodnogo transporta.  
(Aerodynamic models) (Hydraulic models)  
(Rivers)

MARTYNOV, M.I., general-mayor aviatsii, voyenny letchik pervogo  
klassa; FADDEYEV, N.I., polkovnik, voyenny shturman pervogo  
klassa

Isn't it about time we changed the procedure in flight pre-  
paration. Vest.Vozd.Fl. no.1:22-26 Ja '60.  
(MIRA 13:8)

(Flight training)

FADEEVA, N.P. [Fadeyeva, N.P.]; RAUTENSTEIN, I.I. [Rautenshteyn, Ya. I.];  
~~EL'PINER, I.E.~~ [El'piner, I. Ye.]

Influence of ultrasonics on some actinophages and bacteriophages.  
Analele biol 14 no.1:39-45 Ja-Mr '60.

YAKOVLEV, A.A.; FADDEYEV, O.V.

Full-scale test of the icebreaker "I.Stalin" in 1959. Probl.Arkt.1  
Antarkt. no.5:81 '60. (MIR 14:4)

(Ice-breaking vessels)

BESSUDNOV, V.M., inzh.; ~~PADDEYEV~~, O.V., inzh.

Requirements by classification societies of ship strength  
for sailing in ice. Sudostroenie 28 no.1:7-10 Ja '62.

(MIRA 16:7)

(Hulls(Naval architecture))  
(Sea ice)

SINITSYNA, G.S.; PADDNYEV, S.L.; SUKHODOLOV, G.M.

Electrolytic separation of micro quantities of uranium and  
plutonium. Radiokhimiia 1 no.3:295-299 '59. (MIRA 12:10)  
(Uranium) (Plutonium)



~~FADEYEV~~, Sergey Pavlovich [deceased]; ZYBIN, V.P., doktor tekhn. nauk, retsenzent; POKROVSKIY, A.M., kand. tekhn. nauk, dots., nauchn. red.; KOLODYAZHNAYA, Zh.A., red.

[Design of machine parts; collection of problems] Raschety detalei mashin; sbornik zadaoh. Moskva, Vysshaya shkola, 1964. 180 p. (MIRA 18:3)

1. Zaveduyushchiy kafedroy "Detali mashin. PTU" Vsesoyuznogo zaochnogo instituta tekstil'noy i legkoy promyshlennosti (for Zybin).

VORONOVSKIY, V.R.; FADDEYEV, V.P.

Determining the required frequency for transmitting information  
on the yield of oil wells. Nefteprom. delo no.9:21-25 '65.

(MIRA 18:10)

1. Vsesoyuznyy nauchno-issledovatel'skiy i proyektno-konstruktorskiy  
institut kompleksnoy avtomatizatsii neftyanoy i gazovoy promyshlen-  
nosti.

FADDEYEV, Ye.T.,

Fascinating trip. Nauka i zhizn' 23 no,10:37-38 0 '56.(MLRA 9:11)  
(Moscow--Atomic power--Exhibitions)

AUTHOR: Faddeyev, Ye.T.

SOV '25-58-11-18/44

TITLE: ~~The Transformation of Nature (Preobrazovaniye prirody)~~

PERIODICAL: Nauka i zhizn', 1958, Nr 11, pp 41-48 and p 6 of centerfolds (USSR)

ABSTRACT: This is an anti-religious article dealing with the power and ability of men of changing nature by their own effort and will, independent of any religious beliefs or rules established by the Church. Academician A.Ye. Fersman, K.E. Tsiolkovskiy and V.L. Komarov are mentioned in this connection. There are 6 sketches, 3 pictures and 3 Soviet references.

Card 1/1

FADDEYEV, Ya. T., PLATONOV, G.V., doktor filosof.nauk, nauchnyy red.;  
SPIRIDONOVA, O.I., red.

[Science and religion; album] Nauka i religia; al'bom -  
vystavka. Red.G.V.Platonov. Leningrad, Sovetskaya Rossiya,  
1959. 41 l. [Instructions for the use of the album  
"Science and religion."] Metodicheskie ukazaniya k al'bomu  
"Nauka i religia." 7 p. (MIRA 12:10)  
(Science and religion)

BRYUKHANOV, Valentin Andreyevich [deceased]; FADDEYEV, Ye.T., otv.red.;  
VARVAROV, N.A., otv.red.; STEPANYAN, N.I., red.; ROZEN, E.A.,  
tekhn.red.

[Great achievement of mankind; problem of interplanetary flights  
and atheism] Velikii shag chelovechestva; problema mezhplanetnykh  
poletov i ateizm. Moskva, Izd-vo "Sovetskaya Rossiya," 1959.  
98 p. (Interplanetary voyages) (Atheism) (MIRA 13:3)

21/43

S/025/61/000/006/005/007  
D244/D305

27.0000

4112, 1121

AUTHOR:

Faddeyev, Ye. T.

TITLE:

What is this telepathy?

PERIODICAL: Nauka i zhizn', no. 6, 1961, 60-63

TEXT: The author discusses the question of telepathy with special reference to a recent conference on the philosophic aspects of the natural sciences attended by lecturers from Moscow higher educational institutes. The meeting held was under the auspices of the Kafedra dialekticheskogo i istoricheskogo materializma yestiestvennykh fakul'tetov MGU (Department of Dialectical and Historical Materialism in the Natural Sciences Faculty of Moscow State University. The main paper was read by Doctor of Philosophical Sciences A. G. Spirkin who initially cited some facts and experiments on thought-transmission mentioned in Nauka i zhizn', no. 11, 1960; Znaniye-sila, no. 12, 1960; Tekhnika-molodezhi, nos. 1-3, 1961. He then proposed a new concept of the materialistic effect of organisms on each other to explain the

Card 1/4

21743  
S/025/61/000/006/005/007  
D244/D305

What is this telepathy?

nature of this phenomenon which must interest not only philosophers but also biologists, physiologists, doctors, cyberneticians and engineers. Many scientists favored electromagnetic vibrations as the means whereby thought-transmission takes place, although their views were discounted by V. Arkad'yev on the grounds of the feebleness of the biocurrent in the brain. Other speakers, however, suggested the existence of a special physical nerve-field in the brain. In the author's opinion, thought-transmission is accomplished by means of material agents or signals which transport information about specific thought-processes. These then induce in the brain of the recipient a similar condition to that existing in the brain of the transmitter at the moment of emission of the signals. Thus, in contrast to some views propounded in popular-science literature on telepathy (Tekhnika-molodezhi, no 1, 1961), the author maintains that only material processes, whose nature is not known at present, and not sound signals, serve as the means of stimulating mental patterns during telepathic communication.

Card 2/4



21743

S/025/61/000/006/005/007  
D244/D305

What is this telepathy?

The author continues his argument by noting that the telepathic interrelationship of the subject and object is being studied by physiologists, psychologists and gnoseologists who all agree on the existence of the external world, the brain, and the intermediate organ - the human body. The direct influence of external factors (wounds, shock), however, are not considered to be completely characteristic of the system: object - subject. However, recent data indicate the presence of other phenomena - electromagnetic and magnetic radiation fields, which react on the brain while eluding the body. These little understood processes are of much interest in view of man's entry into space where radiation effects will be more diverse than those on Earth. According to some electrophysiologists and biophysicists, the brain is capable of generating radiation with a differing wavelength and may, therefore, be a source of electromagnetic fields. A new scheme has been proposed whereby one brain can react on another by means of a definite material factor - electromagnetic vibrations;

X

Card 3/4

21743

S/025/61/000/006/005/007

D244/D305

What is this telepathy?

the telepathic link may possibly be through the reactions of the external medium on the organ of consciousness. The author, however, does not agree with such a concept; the telepathic link may well be a new material-agent, about which nothing is known at present. Of greater probability is the existence of special nerve fields - a hypothesis connected with the view that field forms of the movement of matter preceded the development of at any rate highly-organized life. These gravity, electron-positron, meson and nucleon fields appeared at a definite pre-biologic and pre-social stage of evolution; such forms are already known at atomic and stellar levels in the microcosm and macrocosm. No new field forms have since appeared, and it is suggested that the life cell (including the nerve cell) may become the source of a new type of field developed at a higher evolutionary stage and which may be the cause of brain-brain reactions. Another possibility is the perfection of this mental-reaction system in a pre-field state. There are 2 figures and 5 Soviet-bloc references.

Card 4/4

FADDEYEV Yu. I.

SOV / 124-58-5-5389

Translation from: Referativnyy zhurnal, Mekhanika, 1958, Nr 5, p 62 (USSR)

AUTHOR: Faddeyev, Yu. I.

TITLE: Using Energy Relationships to Investigate the Rolling of Ships Under Conditions in a Regular Seaway (Primeneniye energeticheskikh sootnosheniy k voprosu issledovaniya bortovoy kachki sudov na regul'yarnom volnenii)

PERIODICAL: Tr. Leningr. korablestroit. in-ta, 1956, Nr 18, pp 159-174

ABSTRACT: The general problem of the side-to-side roll of a ship is examined under conditions of regular sea waviness. This is a further development of the subject of a previous paper by the author (Tr. Leningr. korablestroit. in-ta, 1955, Nr 15, pp 53-61; see also RZhMekh, 1956, Nr 11, abstract 7492). The equation for ship roll is solved by the energy method; taken into account are the various shapes of the static-stability curve, such as linear and nonlinear functions of the angle of heel, and various analytical relationships between the rotary damping moment and the angular rolling velocity. A graphic method is proposed for arriving at the nonlinear resonance amplitude. The author deems that his method yields more precise results

Card 1/2

SOV / 124-58-5-5389

Using Energy Relationships (cont.)

than do other methods; also, the laboriousness of the calculations is reduced. By way of numerical example, a determination is made of the resonance amplitude of a fishing trawler (displacement 936 tons) by means of both the linear and the nonlinear theories. The results obtained in the two cases are compared. Bibliography: 7 references.

V.B. Dragomiretskiy

1. Ship--Roll
2. Mathematics--Applications

Card 2/2

FADDEYEV, Yu. I. Cand Tech Sci -- (diss) "Study of Lateral Pitching  
of Ship Models on Calm and <sup>Turbulent</sup> ~~Heaving~~ Waters." Len, 1957. 23 pp 20 cm.  
(Len ~~XXXXXXXXXXXX~~ Ship-Designing Inst), 140 copies (KL,26-57,109)

VOYTKUNSKIY, Ya.I., kand.tekhn.nauk; KATSMAN, F.M., inzh.; FADDEYEV, Yu.I.,  
kand.tekhn.nauk; YAKONOVSKIY, S.V., inzh.

Towing resistance of lifeboats. Sudostroenie 24 no.12:15-20  
D '58. (MIRA 12:2)

(Lifeboats)

(Towing)

(Ship resistance)

FADDEYEV, Yu.I.

Oscillation of bodies in a liquid. Trudy LKI no.28:73-80 '59.  
(MIRA 15:5)

1. Kafedra gidromekhaniki Leningradskogo kroablestroitel'nogo  
instituta.

(Ships--Hydrodynamics)

16.3000

24.4 35004

S/044/61/000/003/005/014  
C111/C333

AUTHOR: Faddeyev, Yu. I.

TITLE: The construction of the plane potential flow of an incompressible fluid with the method of conformal mappings for the representation of the solution in parameter form

PERIODICAL: Referativnyy zhurnal, Matematika, no. 3, 1961, 31, abstract 3B127. (Tr. Leningr. Korablestroit. in-ta, 1959, vyp 29, 117-126)

TEXT: The author considers an instationary motion free of circulation of the contour. It is assumed that the function  $z = f(\zeta)$ , which maps the exterior of the contour to the exterior of the circle, is known. The author represents the boundary values of the components of the complex potential  $w(\zeta)$ , which correspond to the translation and to the rotary motion of the contour, as conjugate trigonometric series, uses the boundary conditions and expresses the coefficients of the series mentioned by the coefficients of the expansion of the mapping function. The adjoint masses are expressed by the same coefficients. [Abstracter's note: Complete translation.]  
Card 1/1



FADDEYEV, Yu.I.; LU CHEY, student

Geometric and hydrodynamic characteristics of one family of  
simplest symmetrical profiles. Trudy LKI no.34:81-92 '61.  
(MIRA 15:8)

1. Kafedra gidromekhaniki Leningradskogo korablestroitel'nogo  
instituta (for Faddeyev). 2. Korablestroitel'nyy fakul'tet  
Leningradskogo korablestroitel'nogo instituta (for Lu Chey).  
(Blades)

FADDEYEV, Yu.I.

The motion of a flat deformable shape. Trudy LKI no.36:39-48 '62.  
(MIRA 16:12)

1. Kafedra gidromekhaniki Leningradskogo korablestroitel'nogo  
instituta.

FADDEYEV, Yu.I.; YUY SYAN-SAN [Yü Haiang-san]

Determining coefficients of resistance in the case of high  
damping. Trudy LKI no.38:151-155 '62. (MIRA 16:7)

1. Kafedra gidromekhaniki Leningradskogo korablestroitel'nogo  
instituta (for Faddeyev). 2. Kafedra teorii korablya  
Leningradskogo korablestroitel'nogo instituta (for Yuy Syan-San).  
(Vibration (Marine engineering))

LUGOVSKIY, V.V., kand. tekhn. nauk; FADDEYEV, Yu.I., kand. tekhn. nauk

Calculating the rolling of ships using the nomograms of a  
series of model experiments. Sudostroenie 29 no.8:26-30 Ag '63.  
(MIRA 16:10)

(Ships--Hydrodynamics) (Ship models--Testing)

*FADDEYEV, Yu. N.*

KOVALEVSKIY, Georgiy Nikolayevich; FADDEYEV, Yu.N., red.; IVANOV, K.A., red.;  
TROFIMOV, A.V., tekhn.red.

[Ship theory] Teoriia korablia. Moskva, Izd-vo "Morskoi transport,"  
1956. 259 p. (MIRA 11:1)  
(Hulls (Naval architecture))

FADDEYEVA, A.P.

6589

FADDEYEVA, A. P.

FADDEYEVA, A. P. INZHENERNO-GEOLICHESKIYE IZYSKANIYA NA  
MORSKIKH POBEREZH'YAKH I V UST(YAKH REK. METOD. UKAZANIYA  
L.-M. GOS. 120 LIT PO STROITEL'STVU I ARKHITEKTURE, 1954  
136 s. S ILL. 26 SM. ( ILVO STROITEL'STVA SSSR. GLAVSTROIROY-  
ekt. LENINGR. PROEKTINYY IN-t) 1,000 EKZ 7 R. 90 K. --  
AVT. RAZRABOTKI UKAZAN NA 4\*Y S. - BIBLIOGR: s 133-134  
(65 IAZV.)- (552596) P

624.13:55 plus 016.3)

SO: ZNICHALYA LETOPIS' NO. 6. 1955

LAVROVA, M.A., red.; FADDEYEVA, A.P., red.; ZHINGAREVA--  
DOBROSEL'SKIY, A.T., red.; TOKAIEVA, T.N., ved. red.

[Problems of the stratigraphy of Quaternary sediments  
in the northwestern area of the European part of the  
U.S.S.R.] Voprosy stratigrafii chetvertichnykh otlozhenii  
Severo-Zapada Evropeiskoi chasti SSSR; sbornik statei.  
Leningrad, Gostoptekhizdat, 1962. 198 p. (MIRA 18:5)

1. Nauchno-tehnicheskoye gornoye obshchestvo, Moscow.  
Leningradskoye oblastnoye upravleniye.

ACC NR: AP6030864

SOURCE CODE: UR/0365/66/002/005/0576/0580

AUTHOR: Zemskov, G. V.; Kogan, R. L.; Dombrovskaya, Ye. V.; Kostenko, A. V.;  
Shevchenko, I. M.; Kosa, Ye. V.; Fadeyeva, E. V.; Khmelevskaya, M. Ye.; Mikotina, N. F.

ORG: Odessa Polytechnical Institute (Odesskiy politekhnicheskiy institut)

TITLE: Protective diffusion coatings of nickel alloy

SOURCE: Zashchita metallov, v. 2, no. 5, 1966, 576-580

TOPIC TAGS: <sup>alloy</sup>nickel chromium alloy, aluminum containing alloy, titanium containing alloy, tungsten containing alloy, <sup>alloy</sup>protective coating, <sup>alloy</sup>corrosion resistance, diffusion coating alloy, alloy oxidation resistance/ZhS6-K alloy

ABSTRACT: A series of diffusion coatings were tested for protection of ZhS6-K nickel base alloy (0.13-0.20% carbon, 10.5-12.5% chromium, 5-6% aluminum, 2.5-3% titanium, 2.5-3% tungsten, 4.5-5.5% molybdenum, 0.13-0.20% boron) against gas corrosion in a mixture of products of sulfurous fuel combustion and sea water vapors after all attempts to improve alloy oxidation resistance by alloying failed. Alloy specimens were diffusion coated with one or two elements used simultaneously or one after the other. The coating was done by a pack cementation at 900-1000C for 10 hr. Chromium, aluminum, silicon, titanium, boron, cerium, beryllium and magnesium were used as single-element coatings. Chromium with titanium, silicon, aluminum, or boron; aluminum with boron, cerium, or titanium; titanium with silicon or boron; manganese with boron;

UDC: 621.793.4

Card 1/4



ACC NR: AP6030864





Fig. 1. Dependence of the change of the diffusion layer depth upon the content of elements in the mixture

a - Aluminum-silicon impregnation; b - boron-titanium impregnation.

Card 2/4

L 44.777-66

ACC NR: AP6030864



**Fig. 2. Depth of corrosion in coated and uncoated ZhS6-K alloy.**

cerium with boron; and silicon with aluminum were used for binary coatings. Corrosion tests were done in combustion products containing 0.74% and 0.11% sea water at 900C for 15 hr. It was found that all the coatings tested have a higher corrosion resistance than the uncoated alloy (see Fig. 1). Binary coatings protect the alloy more efficiently than single-element coatings, especially with the consecutive method of

Card 3/4

I. 44077-66

ACC NR: AP6030864

application. Coatings obtained by this method have a higher concentration of elements and a more uniform structure of the surface layer than the coatings applied by other methods. Orig. art. has: 5 figures. [ND]

SUB CODE: 11, 13/ SUBM DATE: 13Jul65/ ATD PRESS: 5077

*all in*  
Card 4/4

TROITSKAYA, Mariya Nikolayevna; FADDEYEVA, I.I., red.; LAZAREVA, L.V.,  
tekhn. red.

[Textbook on laboratory work concerning soil mechanics] Poso-  
bie k laboratornym rabotam po mekhanike gruntov. Moskva, Izd-  
vo Mosk. univ., 1961. 303 p. (MIRA 15:1)  
(Soil mechanics—Research)

ZAKHAROVA, Yelena Mikhaylovna; FADDEYEVA, I.I., red.; YERMAKOV, M.S.,  
tekhn. red.

[Sluicing and the analysis of heavy mud residues from sluicing]  
Shlikhovye poiski i analiz shlikhov. Moskva, Izd-vo Mosk. univ.,  
1959. 163 p. (MIRA 14:10)

(Mineralogy, Determinative)

VOSKRESENSKIY, Sergey Sergeyevich; FADDEYEVA, I.I., red.; GEORGIYEVA,  
G.I., tekhn.red.

[Geomorphology of Siberia] Geomorfologiya Sibiri. Moskva, Izd-vo  
Mosk.univ., 1962. 351 p. (MIRA 15:5)  
(Siberia—Geomorphology)

MILANOVSKIY, Yevgeniy Yevgen'yevich; KHAIN, Viktor Yefimovich; MURATOV, M.V.,  
red.; FADDEYEVA, I.I., red.; MUKHINA, L.V., tekhn.red.

[Geology of the Caucasus.] Geologicheskoe stroenie Kavkaza. [Moskva]  
Izd-vo Mosk. univ., 1963. 355 p. (Ocherki regional'noi gelologii SSSR,  
no.8). (MIRA 16:9)

MARKOVICH, Ye.M.; FADDEYEVA, I.Z.

A new middle Jurassic fern from the Orsk brown coal basin.  
Paleont.zhur. no.3:127-130 '60. (MIRA 13:10)

1. Laboratoriya geologii uglya Akademii nauk SSSR.  
(Aktyubinsk Province--Gerns, Fossil)



FADDEYEVA, I.Z.

Jurassic megaspores from western Kazakhstan. Paleont. zhur. no.4:  
125-128 '60. (MIRA 14:1)

1. Laboratoriya geologii uglya AN SSSR.  
(Kazakhstan—Spores (Botany), Fossil)

VOLKOV, V.N.; MARKOVICH, Ye.M.; FADDEYEVA, I.Z.; VOLKOVA, I.B.

Short review of the history of the study of lower Mesozoic sediments  
in the southern Magnitogorsk synclinorium. Trudy Lab.geol.ugl.

no.12:9-14 '61.

(MIRA 14:8)

(Ural Mountains—Coal geology)

FADDEYEVA, Irina Zakharovna; LYUBER, A.A., kand. geol.-miner.  
nauk, otr. red.

[Palynological basis of the stratigraphic division of Lower  
Mesozoic coal-bearing sediments in the Or'-Ilek region] Pa-  
linologicheskoe obosnovanie stratigraficheskogo raschlene-  
niia nizhnemezozoiskikh uglenesnykh otlozhenii Or'-Ilebskogo  
raiona. Moskva, Nauka, 1965. 116 p. (MIRA 18:8)

FADDEYEVA, Tat'yana Dmitriyevna; NECHAYEV, S.V., red.; SENCHILO, K.K.,  
tekh.n.red.

[Cholera and its control] Kholera i bor'ba s nei. Moskva,  
Gos.isd-vo med.lit-ry Medgis, 1959. 231 p.

(MIRA 14:2)

(CHOLERA, ASIATIC--PREVENTION)

FADDEYEVA, M.D.

Spectral characteristics of the DNA - Acridine Orange complex.

Tsitologia, 4, no. 2, 231 Mar-Apr 1962.

(MIRA 15:8)

1. Laboratoriya biokhimii kletki Instituta tsitologii AN SSSR,  
Leningrad.

(NUCLEIC ACIDS) (ACRIDINE ORANGE)

L 8944-66 EWT(1)/EWA(j)/EWA(b)-2 RO

ACC NR: AP5026554

SOURCE CODE: UR/0286/65/000/019/0111/0111

AUTHORS: <sup>55</sup>Baskakov, Yu. A.; <sup>55</sup>Faddeyeva, M. I.; <sup>55</sup>Andreyeva, Ye. I.; <sup>55</sup>Golyshin, N. M.;  
Novikova, R. G. <sup>55</sup>

ORG: none

TITLE: Method for obtaining fungicidal derivatives of N-carboalkoxyarylhydroxyl amines. Class 45, No. 175347 /announced by All-Union Scientific Research Institute for Chemical Agents for Protection of Plants (Vsesoyuznyy nauchno-issledovatel'skiy institut khimicheskikh sredstv zashchity rasteniy)/

SOURCE: Byulleten' izobreteniy i tovarnykh znakov, no. 19, 1965, 111

TOPIC TAGS: <sup>55</sup>fungicide, arylhydroxyl amine, plant disease control.

ABSTRACT: This Author Certificate presents a method for obtaining fungicidal derivatives of N-carboalkoxyarylhydroxyl amines by reacting alkylchlorocarbonates with arylhydroxylamines. To increase the variety of fungicides, halogen arylhydroxylamines are used as arylhydroxylamines.

SUB CODE: 07/ SUBM DATE: 22Jul64

Card 1/1 <sup>06</sup>

UDC: 632.951.2.547  
547.555

FADDEYEV, D.K.; FADDEYEVA, V.I.

M. Parodi's book "Localization of the eigenvalues of matrices and their application". Zhur. vych. mat. i mat. fiz. 1 no.3:551-552  
My-Je '61. (MIRA 14:8)

(Eigenvalues)  
(Parodi, M.)

**FADDEYEVA, V.**

Vera Petrovna Golovina. Zh. vysshei nerv. deiat. 1 no. 5:784 Sept-Oct 1951. (CML 23:3)

1. Obituary.



FADDEYEVA, V.K.

Effect of phenamine on of the higher nervous function in white rats. Zh. vysshei nerv. deiat., Pavlova 1 no. 2:165-186 Mar-Apr 1951. (GLML 22:5)

1. Department of the Pathophysiology and Therapy of Higher Nervous Activity, Institute of Higher Nervous Activity of the Academy of Sciences USSR.

PADDEYEVA, V.K.

Nature of correlation of the first and second signal systems in formation of conditioned reactions to complex stimulus in children. Zh. vysshei nerv. deiat. Pavlova 1 no.3:361-375 May-June 1951. (CJML 23:2)

1. Department of the Pathophysiology and Therapy of Higher Nervous Activity, Institute of Higher Nervous Activity of the Academy of Sciences USSR.

FADDEYEVA, V. K.

Conditional Response

Effect of phenocoll upon the conditional connection between the primary and secondary reflex system., Zhur, vys. nerv. deiat., 1, No. 6, 1951.

9. Monthly List of Russian Accessions, Library of Congress, April 1952, Unclassified.

FADDEYEVA, V. K.

Dissertation: "Experimental Investigations of the Effect of Phenamine on the Work of Higher Divisions of the Central Nervous System (on the Work of the Cerebrum)."  
Dr Med Sci, Inst of Higher Nervous Activity, Acad Sci USSR, Moscow, Oct-Dec 53.  
Vestnik Akademii Nauk, Moscow, Jan 54)  
(Source gives brief summary of work.)

SO: SUM 318, 23 Dec. 1954